

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁷ : A61M 31/00</p>	<p>A1</p>	<p>(11) International Publication Number: WO 00/57946 (43) International Publication Date: 5 October 2000 (05.10.00)</p>
<p>(21) International Application Number: PCT/US00/07622 (22) International Filing Date: 21 March 2000 (21.03.00) (30) Priority Data: 09/277,359 26 March 1999 (26.03.99) US (71) Applicant: TRANSON, LLC [-/US]; 650 California Street, 25th Floor, San Francisco, CA 94108 (US). (72) Inventors: BOND, Geoffrey; 20 E. Lake Street, Lakewood, NY 14750 (US). PETERSON, Thomas, M.; 4249 Neptune Street, Erie, PA 16506 (US). (74) Agent: HACKLER, Walter, A.; 2372 S.E. Bristol, Suite B, Newport Beach, CA 92660 (US).</p>		<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published With international search report. With amended claims and statement.</p>
<p>(54) Title: METHOD AND APPARATUS FOR EMERGENCY TREATMENT OF A PATIENT EXPERIENCING A THROMBOTIC VASCULAR OCCLUSION</p>		
<p>(57) Abstract</p> <p>This invention is an acute care method, and apparatus (10) for treating a patient experiencing a thrombotic vascular occlusion that includes introducing a selected dose of an active agent proximate a vascular occlusion in the patient in order to lyse the vascular occlusion, radiating the vascular occlusion, and active agent.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

WO 00/57946

PCT/US00/07622

METHOD AND APPARATUS FOR EMERGENCY TREATMENT
OF PATIENTS EXPERIENCING A THROMBOTIC
VASCULAR OCCLUSION

5 The present invention is generally related to
a method and apparatus for pre-hospital or initial
treatment of patients experiencing a thrombotic
vascular occlusion, and is more particularly
10 directed to an emergency application of ultrasound
and an agent for lysing vascular occlusions or
thrombi. Importantly, the agent may not have
activity for lysing vascular occlusions without
application of ultrasound.

15 Thrombosis can cause partial or total occlusion
of blood vessels which leads to a number of
important cardiovascular complications, including
unstable angina, acute myocardial infarction (heart
attack), cerebral vascular accidents (stroke)
20 pulmonary embolism, deep vein thrombosis and
arterial thrombosis.

 It is well known that acute myocardial
infarction is one of the greatest causes of death in
25 the United States, and it has been recognized that
time is of the essence in successfully treating
individuals undergoing acute myocardial infarction.
Many hospitals and caregiving institutions have
established coronary care units with trained
30 personnel and equipment for treating a patient in
the shortest time possible upon arrival. However,
important time is lost during the delivery of
patients to hospitals.

35 Removal or lysing of vascular occlusions, or
clots, may be accomplished through a variety of

WO 00/57946

PCT/US00/07622

agents such as, for example, urokinase, streptokinase, aspirin and tissue plasminogen activators, tPA. These clot dissolving agents are generally used in order to lyse clots which have
5 formed in the coronary arteries.

These agents are typically injected into the bloodstream or organ close to the position of the clot. Unfortunately, such agents have a side effect
10 of causing undesirable bleeding in the patient. Thus, patients who may have an ulcer or other bleeding disorder are especially difficult to treat with systemic anticoagulants. Aspirin has found
15 wide use as a clot inhibitor, particularly with respect to clots on the arterial side of circulation. However, aspirin is used as an adjuvant agent to prevent thrombosis and has little if any effect on an established blood clot.

20 Ultrasound has found use in the dissolution of vascular occlusions. For example, U.S. Patent Nos. 5,269,291, 5,318,014, 5,362,309, 5,431,662 and 5,474,531, describe intravascular ultrasonic tools for the dissolution of intravascular blockages.

25 More recently, transcutaneous ultrasound has been found to enhance the activity of thrombolytic agent, for example, see U.S. Patent Nos. 5,509,896 and 5,695,460. All of the hereinabove referenced
30 patents have been assigned to the assignee of the present application.

Thus, while it has been recognized that the use of ultrasound with an active agent enhances the
35 activity of the agent in lysing of a vascular occlusion, other synergistic properties have not been heretofore discovered. For instance, it has

WO 00/57946

PCT/US00/07622

been found in accordance with the present invention that ultrasound shortens the onset of thrombolysis activity of an agent such as, for example, microbubbles or tPA. Accordingly, more rapid onset
5 of dissolution of clots results in less damage from the vascular occlusion because of more rapid opening of the blood vessel and in the restoration of tissue perfusion. In addition, the combination of certain agents and ultrasound could reduce the likelihood of
10 excess bleeding which may significantly increase the likelihood of survival by a patient.

SUMMARY OF THE INVENTION

15 A method in accordance with the present invention for providing acute care treatment of a patient experiencing thrombotic vascular occlusion generally includes the steps of introducing a selected dose of an agent for acting on a vascular
20 occlusion in the patient in order to lyse the vascular occlusion and irradiating the vascular occlusion and active agent in order to shorten onset and accelerate the effectiveness of lysing action of the agent. In view of the well known urgency of
25 heart attack and stroke matters, i.e., cell death is directly proportional to time, it is of utmost importance to enhance the onset and accelerate the effectiveness of the active agent in lysing the vascular occlusion.

30

An "active agent" as used herein is meant to include an agent having little or no lysing activity without ultrasound, but exhibiting lysing activity with ultrasound.

35

The method in accordance with the present invention applies to coronary, cerebral and

WO 00/57946

PCT/US00/07622

peripheral vascular (venous or arterial) occlusions and it is found that the active agent may be a thrombolytic agent, anticoagulants, agents which alter blood viscosity and supply nuclei to facilitate microstreaming, cavitation agents that enhance clot disruption (e.g., Hespan, Pentaspan, an intravenous fat emulsion such as intralipid, liposyn, a microbubble medium, and an antiplatelet agent or other suitable lysing agents).

10

Accordingly, apparatus in accordance with the present invention, includes a self-contained mobile unit for paramedic or emergency treatment of a patient experiencing thrombotic vascular occlusion. In view of the utmost importance of availability, the apparatus includes a plurality of active agents for dissolution of vascular occlusions. A tutored paramedic or physician selects an agent based upon the indication of patient bleeding and intravenously supplies the agent to the vascular occlusion along with transcutaneously radiating the vascular occlusion.

15
20

Accordingly, one method in accordance with the present invention includes application of ultrasound transcutaneously. Under certain situations, the ultrasound may be applied intravascularly and the active agent may be introduced proximate the vascular occlusion by direct injection.

25
30

BRIEF DESCRIPTION OF THE DRAWINGS

The advantages and features of the present invention will be better understood by the following description when considered in conjunction with the accompanying drawings, in which:

35

WO 00/57946

PCT/US00/07622

Figure 1 is a diagram of apparatus in accordance with the present invention which also illustrates the method in accordance with the present invention; and

5

Figure 2 is a graph of lysing effectiveness with and without ultrasound as a function of time.

DETAILED DESCRIPTION

10

With reference now to Figure 1, there is shown apparatus 10 for effecting pre-hospital or emergency treatment of a patient experiencing thrombolytic vascular occlusion which includes a self-contained mobile unit 12. As represented by boxes 14, 16, 18, 20, 22, 24, a plurality of active agents for dissolution of vascular occlusion are provided with at least one of the active agents being selected by a paramedic or physician for intravenous introduction into a patient 30 by a vial 32, valve 34 and line 36 which, in combination, provide intravenous means for introducing the selected agent into the patient 30. A vascular occlusion in the present application includes coronary occlusions, cerebral occlusions and peripheral occlusions both venous and arterial. A portable ultrasonic oscillator 40 and transducer 42, in combination, provide a means for transcutaneously radiating a vascular occlusion as indicated by the dotted line at 46. The oscillator 40 may be battery powered, connected to a vehicle electrical system, or connected to a conventional electrical source (not shown) by any conventional means. It should be appreciated that multiple oscillators and transducers (not shown) may be utilized, each having a desirable operating range and properties.

WO 00/57946

PCT/US00/07622

It should be appreciated that ultrasound may be applied intravascularly under proper circumstances, and, in this instance, a miniature transducer, and connected to the oscillator 40 as indicated by the dotted line 50 may be utilized. In this regard, reference is made to U.S. Patent No. 5,269,291, 5,318,014, 2,062,309, 3,431,662, and 4,474,531 which are hereby incorporated in toto by this specific reference thereto as indicating intravascular transducers which may be utilized in accordance with the present invention, all of these patents being assigned to assignee of the present invention.

The transducer 42 may be of any suitable design, for example, as set forth in U.S. Patent No. 5,879,314 which is incorporated herewith in toto by this reference thereto for the purpose of illustrating the types of transcutaneous transducers which may be suitable for use in accordance with the present invention. The effective frequency range may be from between about 10 kHz to about 2 MHz, with a desirable frequency range being between about 20 kHz and about 500 kHz and another desirable frequency range being between 20 kHz and 50 kHz. The oscillator 40 may be of any suitable type as set forth in any number of the hereinabove referenced U.S. patents.

As hereinabove noted, timing and selection of appropriate lysing agents is of utmost importance.

For example, a criteria for selection between an echo contrast agent such as sonicated albumin, perfluorocarbon, a lysing or thrombolytic agent such as streptokinase or tPA, an anticoagulant such as heparin, antiplatelet agents, such as platelet

WO 00/57946

PCT/US00/07622

receptor, like a GP IIb-IIIa inhibitor, such as blockers, Aggrastat, Integrillin, a GP IIb-IIIa platelet inhibitor, Reopro, a hyperalimentation agent such as intravenous fat emulsions, intralipid and liposyn, or another alternative agent such as Hetastarch or other artificial colloidal solutions, such as Pentaspan®, may be as follows.

If there is no indicated bleeding, or significant risk of bleeding, a thrombolytic agent may be selected. If there is possible indicated or risk of bleeding, a GP IIb-IIIa inhibitor may be selected. If there is an absolute contraindication due to risk of bleeding, a microbubble, Hetastarch or Pentaspan is preferably selected.

Other indications may be utilized for agent selection.

It should also be appreciated that a combination of agents may be utilized, for example, an echocontrast agent may be used in combination with a thrombolytic agent. For example, an echocontrast agent may be a perfluorocarbon, such as, for example, the dodecafluoropentane colloid dispersion. The echocontrast agent may be a microbubble medium such as free gas bubbles, stabilized gas bubbles, colloidal suspensions, emulsions and aqueous solutions other than dodecafluoropentane. A thrombolytic agent may be any agent having suitable activity such as, for example, streptokinase, staphlokinase, urokinase or a tissue plasminogen activator (tPA). These agents are set forth herein only by way of example and it should be appreciated that, as hereinabove recited, any thrombolytic agent has possible use in accordance with the present invention.

WO 00/57946

PCT/US00/07622

As hereinabove noted, a Hetastarch such as HESPAN®, which is a plasma volume expander, has been found to be effective when used in combination with ultrasound for the lysing of vascular occlusions.

5

Hetastarch is an artificial colloid derived from a waxy starch composed almost entirely of amylopectin. Hydroxyethyl ether groups are introduced into the glucose units of the starch and the resultant material is hydrolyzed to yield a product with a molecular weight suitable for use as a plasma volume expander and erythrocyte sedimenting agent.

10

15

As also hereinabove noted, a colloid suspension or pentastarch, such as Pentaspan®. Pentaspan® or Pentastarch is an artificial colloid derived from a waxy starch composed almost entirely of amylopectin. Hydroxyethyl ether groups are introduced into the glucose units of the starch and the resultant material is hydrolyzed to yield a product with a molecular weight suitable for use in an erythrocyte sedimenting agent.

20

25

Additionally, the radiation by ultrasound may include continuous or pulse radiation, still more particularly by way of example only, the amount of active agent introduced may be in concentration less than about 2000 microliters.

30

35

The combination of ultrasound has the unique effect of accelerating the onset of lysing activity of a GP IIb-IIIa antiplatelet inhibitor such as Reopro or a GP IIb-IIIa blocker such as, for example, Aggrastat and Integrillin.

WO 00/57946

PCT/US00/07622

Figure 2 shows a percent of lysing of human blood clots in vitro when blood clots were incubated with Aggrastat alone at the blood concentration achieved in patients being treated with this agent alone at a concentration which is similar to that given therapeutically to patients and in combination with ultrasound at a frequency of 20 kHz applied transcutaneously. It is evident that in five minutes the use of ultrasound more than doubles the lysing activity of the Aggrastat with the overall lysing activity of being less enhanced at, for example, 30 minutes. This onset of lysing activity is important in acute care situations.

Further, the results utilizing ultrasound and tPA and PESDA, a sonicated albumin perfluorocarbon are shown in Tables 1 and 2. The data shown in Tables 1 and 2 on lysing of clots formed by electric induction of thrombotic occlusion of the left anterior descending coronary artery (LAD) to test the effects of PESDA and tPA in combination with transcutaneous ultrasound on reperfusion in a dog model in vivo. As indicated in the Tables, the ultrasound frequencies utilized were 26 kHz and 37 kHz. Heparin was also used in combination and the flow of blood through the arteries indicated as TIMI with a TIMI value of 0 representing no flow and a TIMI value of 3 representing unoccluded flow of blood. Other agents such as antiplatelet agents, such as GP IIb-IIIa inhibitor, microbubble mediums, anti-coagulants volume expanders and agents having a particulate nature such as Hetastarch, Pentaspan, as well as anticoagulants such as Heparin, produce similar results in vitro.

35

WO 00/57946

PCT/US00/07622

Tables 1 and 2 show data on ultrasound + PESDA and TPA + USD or TPA alone in a dog myocardial infarction model.

5 The results shown in Tables 1 and 2 indicate that PESDA which has no thrombolytic effect of its own facilitates ultrasound clot lysis in a coronary artery with there being re-establishment of blood flow in 5 of 6 dogs studied after the combination of
10 intravenous PESDA and transcutaneous ultrasound over the dog's chest.

 Table 2 shows that in comparison with intravenous tPA, (given in standard doses for
15 humans) the standard thrombolytic drug given to patients with a heart attack, tPA only dissolves the clot in one of four dogs. Even in this case, there was residual clot in the coronary artery. However, when transcutaneous ultrasound over the dog's chest
20 wall, aimed at the heart, is combined with the intravenous tPA, the clots are all dissolved, and there are no significant residual filling defects from blood clots. These data indicate that ultrasound plus a combination of a drug or drugs can
25 be used to more effectively dissolve blood clots in the arteries of the heart to more effectively treat heart attacks.

 Also incorporated into the present invention is
30 a method, and corresponding apparatus 10, for treating a patient on a non-acute basis using the ultrasound methods hereinbefore displaced with agents which have no separate lysing activity but exhibit, or are activated, lysing activity when used
35 in combination with ultrasound. As hereinabove noted, such agents include plasma volume expanders

WO 00/57946

PCT/US00/07622

Comparison of reperfusion using t-PA vs t-PA plus transcutaneous ultrasound (26 kHz) in a dog model of acute myocardial infarction, electric induction of a thrombotic occlusion of the left anterior descending coronary artery (LAD)						
No	1	2	3	4	5	6
Date	12/24/98	12/31/98	1/15/99	1/19/99	1/21/99	1/28/99
ID	#B-B4	#B-B6	#B-B7	#B-B8	#B-B9	#B-B10
Vessel	LAD	LAD	LAD	LAD	LAD	LAD
Method	t-PA alone	t-PA+USO	t-PA alone	t-PA+USO	t-PA+USO	t-PA+USO
Electricity time	100 min	45 min	200 min	100 min	100 min	160 min
Clot age	60 min	90 min	120 min	120 min	120 min	360 min
Heparin	TIMI 0:2000 IV TIMI 3: 4000 SC	TIMI 0:1000 IV TIMI 3: 2000 SC	TIMI 0:1000 IV	TIMI 0:1000 IV TIMI 3: 2000 SC	TIMI 0:1000 IV TIMI 3: 1000 SC	TIMI 0:1000 IV TIMI 1: 1000 SC
TIMI during treatment	TIMI 3	TIMI 3/15 min	TIMI 0	TIMI 1	TIMI 3 with residual T1 and distal occlusions	TIMI 0
20 min						
40 min	N/A	N/A	TIMI 0	TIMI 1	TIMI 0	TIMI 1
60 min	N/A	N/A	TIMI 0	TIMI 1	TIMI 0	TIMI 1
90 min	N/A	N/A	TIMI 0	TIMI 1	TIMI 0	TIMI 2
TIMI during observation						
30 min	TIMI 3	TIMI 3	N/A	TIMI 1	TIMI 3	TIMI 3
60 min	TIMI 3	TIMI 3	N/A	TIMI 1	TIMI 3	TIMI 3
90 min	TIMI 3	TIMI 3	N/A	TIMI 3	TIMI 3	TIMI 3
90 min angiogram	patent with filling defects	widely patent	LAD occlusion	widely patent	LAD occlusion	patent with minor filling defects
Serum analysis			Yes	Yes	Yes	Yes

TABLE 2

WO 00/57946

PCT/US00/07622

and colloid suspensions such as HESPAN® and PENTASPAN®.

5 Although there has been hereinabove described
a specific arrangement of ultrasonic apparatus and
method for thrombi dissolution in accordance with
the present invention for the purpose of
illustrating the manner in which the invention may
be used to advantage, it should be appreciated that
10 the invention is not limited thereto. Accordingly,
any and all modifications, variations, or equivalent
arrangements which may occur to those skilled in the
art, should be considered to be within the scope of
the present invention as defined in the appended
15 claims.

WO 00/57946

PCT/US00/07622

WHAT IS CLAIMED IS:

5 1. Apparatus for effecting emergency treatment of a patient experiencing a thrombotic vascular occlusion, said apparatus comprising:

 a plurality of active agents for dissolution of vascular occlusions, use of at least one of the active agents being selected on a basis of bleeding
10 indication of the patient;

 means for intravenously introducing a select agent into the patient; and

 portable ultrasonic means of
15 transcutaneously radiating the vascular occlusion.

 2. The apparatus according to claim 1 wherein said plurality of active agents include at least one agent for use with a patient having no indicated bleeding, at least one agent for use with
20 a patient having possible indicated bleeding and at least one agent for use with a patient having indicated bleeding.

 3. A self-contained mobile unit for emergency
25 treatment of a patient experiencing a thrombotic vascular occlusion, the unit comprising:

 a plurality of active agents for dissolution of vascular occlusion, use of at least one of the active agents being selected on a basis of bleeding
30 indication of the patient;

 intravenous means for introducing a selected agent into the patient;

 portable ultrasonic means for
35 transcutaneously radiating the vascular occlusion;
 and

 module means for storing the active agents, intravenous means and ultrasonic means.

WO 00/57946

PCT/US00/07622

4. The apparatus according to claim 3 wherein said plurality of active agents includes at least one agent for use with a patient having not indicated bleeding, at least one agent for use with a patient having possible indicated bleeding and at least one agent for use with a patient having indicated bleeding.
5. The apparatus according to claim 1 or 3 wherein at least one of the active agents is a microbubble medium.
6. The apparatus according to claim 5 wherein the micorbubble medium is comprised of a medium selected from the group comprising free gas bubbles, stabilized gas bubbles, colloidal suspension, emulsion and aqueous solutions.
7. The apparatus according to claim 1 or 3 wherein at least one of the active agents in a thrombotic agent.
8. The apparatus according to claim 1 or 3 wherein at least one of the active agents is an antiplatelet agent.
9. The apparatus according to claim 1 or 3 wherein at least one of the active agents is an anticoagulant.
10. The apparatus according to claim 1 or 3 wherein at least one of the active agent is a blood composition altering agent.
11. The apparatus according to claim 10 wherein said blood composition altering agents are selected

WO 00/57946

PCT/US00/07622

from a group consisting of a blood volume expansion agent, a blood replacement agent, a Hetastarch, Pentastarch and Dextran.

5 12. The apparatus according to claim 1 or 3 wherein at least one of the active agents in a intravenous fat emulsion.

10 13. The apparatus according to claim 12 wherein said intravenous fat emulsion is selected from a group comprising intralipid and liposyn.

15 14. The apparatus according to claim 1 or 3 wherein at least one of the active agents in a GP IIb-IIIa platelet inhibitor.

15 15. The apparatus according to claim 14 wherein the GP IIb-IIIa platelet inhibitor is Reopro.

20 16. The apparatus according to claim 1 or 3 wherein at least one of the active agents is a GP IIb-IIIa platelet blocker.

25 17. The method according to claim 16 wherein the GP IIb-IIIa platelet blocker is selected from a group comprising Aggrastat and Integrillin.

WO 00/57956

PCT/US00/08060

AMENDED CLAIMS

[received by the International Bureau on 26 June 2000 (26.06.00);
original claims 1-17; replaced by new claims 1-15 (3 pages)]

1. Apparatus for effecting emergency treatment
5 of a patient experiencing a thrombotic vascular
occlusion, said apparatus comprising:

a plurality of active agents for dissolution
of vascular occlusions, use of at least one of the
active agents being selected on a basis of bleeding
10 indication of the patient, said plurality of active
agents including at least one agent for use with a
patient having no indicated bleeding, at least one
agent for use with a patient having possible
indicated bleeding and at least one agent for use
15 with a patient having indicated bleeding;

means for intravenously introducing the
selected agent into the patient; and

portable ultrasonic means of
transcutaneously radiating the vascular occlusion.
20

2. A self-contained mobile unit for emergency
treatment of a patient experiencing a thrombotic
vascular occlusion, the unit comprising:

a plurality of active agents for dissolution
25 of vascular occlusion, use of at least one of the
active agents being selected on a basis of bleeding
indication of the patient said plurality of active
agents including at least one agent for use with a
patient having not indicated bleeding, at least one
30 agent for use with a patient having possible
indicated bleeding and at least one agent for use
with a patient having indicated bleeding;

intravenous means for introducing a selected
agent into the patient;

35 portable ultrasonic means for
transcutaneously radiating the vascular occlusion;
and

WO 00/57956

PCT/US00/08060

module means for storing the active agents,
intravenous means and ultrasonic means.

5 3. The apparatus according to claim 1 or 2
 wherein at least one of the active agents is a
 microbubble medium.

10 4. The apparatus according to claim 3 wherein
 the micorbubble medium is comprised of a medium
 selected from the group comprising free gas
 bubbles, stabilized gas bubbles, colloidal
 suspension, emulsion and aqueous solutions.

15 5. The apparatus according to claim 1 or 2
 wherein at least one of the active agents in a
 thrombotic agent.

20 6. The apparatus according to claim 1 or 2
 wherein at least one of the active agents is an
 antiplatelet agent.

25 7. The apparatus according to claim 1 or 2
 wherein at least one of the active agents is an
 anticoagulant.

 8. The apparatus according to claim 1 or 2
 wherein at least one of the active agent is a blood
 composition altering agent.

30 9. The apparatus according to claim 18 wherein
 said blood composition altering agents are selected
 from a group consisting of a blood volume expansion
 agent, a blood replacement agent, a Hetastarch,
 Pentastarch and Dextran.

35

WO 00/57956

PCT/US00/08060

10. The apparatus according to claim 1 or 2 wherein at least one of the active agents in a intravenous fat emulsion.

5 11. The apparatus according to claim 10 wherein said intravenous fat emulsion is selected from a group comprising intralipid and liposyn.

10 12. The apparatus according to claim 1 or 2 wherein at least one of the active agents in a GP IIB-IIIa platelet inhibitor.

15 13. The apparatus according to claim 12 wherein the GP IIB-IIIa platelet inhibitor is Reopro.

14. The apparatus according to claim 1 or 2 wherein at least one of the active agents is a GP IIB-IIIa platelet blocker.

20 15. The method according to claim 14 wherein the GP IIB-IIIa platelet blocker is selected from a group comprising Aggrastat and Integrillin.

WO 00/57956

PCT/US00/08060

STATEMENT UNDER ARTICLE 19 (1)



INTERNATIONAL BUREAU OF WIPO
34, Chemin des Colombettes
1211 Geneva 20
SWITZERLAND
Gentlemen:

In response to an International Search Report, mailed 9 June 2000, the Applicant has limited the scope of the claims to the apparatus of the invention. No new matter is added by this amendment as the new claims correspond to originally filed claims as indicated in the letter accompanying the replacement pages. The present amendment replacing claims does not affect the original description or the original drawings.

WO 00/57946

1/1

PCT/US00/07622

*FIG. 1.**FIG. 2.*

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/07622

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61M 31/00

US CL : 604/500

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 604/500

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EAST

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5,695,460 A [SIEGAL et al.] 19 December 1997, entire document.	1, 3, 5-7
X	US 5,380,273 A [DUBRUL et al.] 10 January 1995, entire document.	1, 3
Y	US 5,498,238 A [SHAPLAND et al.] 12 March 1996, entire document.	8-10
A,P	US 5,961,483 A [SAGE et al.] 05 October 1999, entire document.	14, 16
A,P	US 5,986,065 A [WONG et al.] 16 November 1999, entire document.	15
Y	US 5,498,421 A [GRINSTAFF et al.] 12 March 1996, entire document.	10-13

☐ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

11 MAY 2000

Date of mailing of the international search report

09 JUN 2000

 Name and mailing address of the ISA/US
 Commissioner of Patents and Trademarks
 Box PCT
 Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

CATHERINE SERKE

Telephone No. (703) 308-4846